



SHINE A LIGHT ON AN hATTR-PN DIAGNOSIS

hATTR-PN can seem like other diseases and take years to get the right diagnosis. If a genetic test reveals it's hATTR-PN, **ask your doctor if WAINUA is right for you.**

Actor portrayal.

WAINUA is a prescription medication for the treatment of polyneuropathy caused by hereditary transthyretin-mediated amyloidosis (hATTR-PN) in adults.

Individual results may vary.

IMPORTANT SAFETY INFORMATION

WAINUA may cause side effects, including:

- **Low vitamin A level**, which is a serious, but common side effect of treatment with WAINUA.

See pages 4 and 5 for additional information.

Please see additional Important Safety Information on pages 4 and 5 and full Prescribing Information, including Patient Information, for WAINUA.



WAINUA[®]
(eplontersen) _{45 mg}
injection for subcutaneous use

What is hATTR-PN?

hATTR-PN stands for...

hereditary
(passed down from a parent)

Amyloidosis
(group of diseases caused by misfolded proteins) of

Transthyretin
(protein that transports)

Thyroxine
(thyroid hormone) and

Retinol
(vitamin A)

Misfolded transthyretin (TTR) proteins can result in amyloid buildup and cause damage throughout the body.

Poly
(many)

Neuropathy
(damage to nerves throughout the body)

WAINUA reduces TTR protein production, which results in fewer amyloid deposits throughout your body.

IMPORTANT SAFETY INFORMATION

WAINUA may cause side effects, including:

• **Low vitamin A level**, which is a serious, but common side effect of treatment with WAINUA. Your doctor should tell you to take vitamin A supplements. **Do not** take more than the amount of vitamin A your doctor has recommended. Call your doctor if you develop eye problems such as difficulty seeing at night or in low-lit areas (night blindness), or dry eyes. If you develop eye problems, your doctor should refer you to an eye doctor

Polyneuropathy due to hereditary ATTR amyloidosis can significantly affect your quality of life by causing:

- a range of serious symptoms
- a loss of independence when dressing, bathing, or using the bathroom
- an inability to perform activities of daily living

In hATTR-PN, amyloid deposits most commonly accumulate (build up) in:



The nervous system



The digestive system



The heart

WAINUA is not approved to treat the symptoms of cardiomyopathy.*

Symptoms of hATTR can seem unrelated, as amyloid can deposit in various tissues and organs, and present as more common conditions, making it difficult to diagnose.

Symptoms may vary and can include:

- diarrhea or constipation
- pain in the hands and feet
- muscle weakness, causing difficulty walking

*A disease of the heart that makes it harder for your heart to pump blood.

Please see additional
Important Safety Information
on pages 4 and 5.

 **WAINUA**
(eplontersen)^{45 mg}
injection for subcutaneous use

Shine a light on hATTR-PN with a genetic test

Here's why genetic testing is important:



Genetic testing is used to check for hereditary diseases like hereditary ATTR (hATTR) and is recommended for all patients diagnosed with ATTR amyloidosis, for patients with symptoms consistent with hATTR, and for those with a family history of hATTR.



If you have these symptoms OR have already been diagnosed with ATTR, a **TTR gene test can determine if this is due to hereditary ATTR amyloidosis.**



The test looks at the **TTR gene** to see if any changes are present that could be the cause of your amyloidosis.



The process is simple: a small sample of blood, saliva, or cheek cells is all that's needed. Results usually come back within a few weeks.



Most importantly, detecting a **TTR gene variant** will provide a better understanding of your amyloidosis and **shed light on hATTR risk for others in your immediate family.**

IMPORTANT SAFETY INFORMATION

WAINUA may cause side effects, including:

- **Low vitamin A level**, which is a serious, but common side effect of treatment with WAINUA. Your doctor should tell you to take vitamin A supplements. **Do not** take more than the amount of vitamin A your doctor has recommended. Call your doctor if you develop eye problems such as difficulty seeing at night or in low-lit areas (night blindness), or dry eyes. If you develop eye problems, your doctor should refer you to an eye doctor.

The most common side effects include decreased vitamin A and vomiting.

Before you take WAINUA, tell your healthcare provider:

- if you are pregnant or breastfeeding, or plan to become pregnant or breastfeed. It is not known if WAINUA can harm your baby. Changes in vitamin A levels and vitamin A supplementation related to use of WAINUA may harm your unborn baby. It is not known if WAINUA passes into breast milk



Anna T.

TIME TO hATTR-PN DIAGNOSIS:
7 YEARS

Actor portrayal.

Track your symptoms and start the conversation with your doctor today; it may bring you closer to the clarity and answers you're looking for.

Get the symptom tracker at www.wainua.com

Once your doctor has confirmed you have polyneuropathy caused by hATTR as an adult, ask them if WAINUA should be part of your treatment plan.

IMPORTANT SAFETY INFORMATION (continued)

- **about all the medicines you take**, including prescription and nonprescription medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take vitamin A or beta-carotene supplements

You may report side effects related to AstraZeneca products.

Please see full [Prescribing Information](#), including [Patient Information](#), for WAINUA.

 **WAINUA**[®]
(eplontersen) 45 mg
injection for subcutaneous use

WAINUA works by reducing TTR protein production

hATTR starts with changes in the TTR gene, leading to amyloid buildup that may cause symptoms over time.

The transthyretin protein (TTR) carries vitamin A and thyroid hormones throughout the body. In people with hATTR, the unstable TTR protein breaks apart, misfolds, and forms amyloid fibrils that can build up and cause damage throughout the body.



WAINUA works within the liver, where TTR protein is made,



to lower the production of TTR,



which results in fewer amyloid deposits throughout your body.

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects include decreased vitamin A and vomiting.



“Living with hATTR-PN has been a challenge, but WAINUA has definitely made it a lot more manageable.”

– Tom, actual WAINUA patient
Your results may vary.

Check out how Tom keeps it simple with WAINUA at www.wainua.com/patient-voices.

Just one dose each month gives Tom more time to do activities with his sons. For him, the convenience of self-injection is what matters most.

Please see additional Important Safety Information on pages 4 and 5.

 **WAINUA**[®]
(eplontersen)^{45 mg}
injection for subcutaneous use

Ask your doctor if WAINUA could be your way



WAINUA can prevent your neuropathy from getting worse so you may feel stronger.*



WAINUA can improve quality of life, allowing you to do more of the things you want to do.†

Individual results may vary.

The WAINUA-treated group had a significant improvement in neuropathy impairment (mNIS+7 score) and quality of life (Norfolk QoL-DN score) after 35 weeks in a clinical trial of adults with hATTR-PN.[†]

*As measured by mNIS+7 score and Norfolk QoL-DN score.

†Improvement was defined as the difference between the change from baseline in the WAINUA group and a placebo group from a similar study.

The mNIS+7 scale assessed polyneuropathy by measuring things such as muscle strength, sensation, and reflexes.

The Norfolk QoL-DN questionnaire assessed patients' perceptions of their neuropathy symptoms and its impact on daily activities, such as dressing, bathing, or walking.

IMPORTANT SAFETY INFORMATION (continued)

Before you take WAINUA, tell your healthcare provider:

- **if you are pregnant or breastfeeding, or plan to become pregnant or breastfeed.** It is not known if WAINUA can harm your baby. Changes in vitamin A levels and vitamin A supplementation related to use of WAINUA may harm your unborn baby. It is not known if WAINUA passes into breast milk
- **about all the medicines you take**, including prescription and nonprescription medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take vitamin A or beta-carotene supplements



Safety & Side Effects



WAINUA was generally well tolerated in a clinical trial.



WAINUA suppresses the production of the protein that transports vitamin A and, therefore, vitamin A supplementation is recommended.



No additional laboratory monitoring is needed when you take WAINUA.

IMPORTANT SAFETY INFORMATION

WAINUA may cause side effects, including:

- **Low vitamin A level**, which is a serious, but common side effect of treatment with WAINUA. Your doctor should tell you to take vitamin A supplements. **Do not** take more than the amount of vitamin A your doctor has recommended. Call your doctor if you develop eye problems such as difficulty seeing at night or in low-lit areas (night blindness), or dry eyes. If you develop eye problems, your doctor should refer you to an eye doctor

Please see additional
Important Safety Information
on pages 4 and 5.

 **WAINUA**
(eplontersen)^{45 mg}
injection for subcutaneous use

Take control

WAINUA is the only treatment for hATTR-PN that you can self-administer at home... or wherever life takes you.*



You can take WAINUA at home or on the go

Just make sure you store it properly.*



It's taken once monthly

WAINUA won't wear off between doses.



The actual injection takes just 10 seconds

After preparation,[†] you can take WAINUA by yourself or someone can help you. Be sure to read the complete Instructions for Use.



Easy storage

WAINUA can be stored at room temperature for up to 6 weeks. Otherwise it should be stored in its carton in the refrigerator.*

*Store WAINUA in its carton in the refrigerator between 36 °F to 46 °F (2 °C to 8 °C). Do not freeze. WAINUA can be stored at room temperature up to 86 °F (30 °C) for up to 6 weeks. Do not let WAINUA reach temperatures above 86 °F (30 °C) or expose to heat. Throw away WAINUA if kept at room temperature longer than 6 weeks.

[†]Preparation: bring device to room temperature (30 minutes), check device, and prepare injection site. Read Instructions for Use for full directions. Your healthcare provider should show you how to prepare and inject WAINUA for the first time.

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects include decreased vitamin A and vomiting.

Before you take WAINUA, tell your healthcare provider:

· **if you are pregnant or breastfeeding, or plan to become pregnant or breastfeed.** It is not known if WAINUA can harm your baby. Changes in vitamin A levels and vitamin A supplementation related to use of WAINUA may harm your unborn baby. It is not known if WAINUA passes into breast milk



Paul S.

TIME TO hATTR-PN DIAGNOSIS:
4 YEARS

Actor portrayal.

“With WAINUA, I can take my dog on a long walk... I’m able to instruct my fitness classes. It’s been wonderful.”

– Brigitte, actual WAINUA patient
Your results may vary.

Check out real patients taking WAINUA at www.wainua.com/patient-voices. It perfectly fits their structured lifestyles, giving them the freedom to manage their health on their own terms.

Not sure what to ask at your next doctor appointment?

For questions to ask and space to take notes, download and print the doctor discussion guide and then bring it to your next appointment.

To download the discussion guide

visit WAINUA.com/discussion-guide

Please see additional
Important Safety Information
on pages 4 and 5.

 **WAINUA**
(eplontersen)^{45 mg}
injection for subcutaneous use

WAINUA^{Way} is committed to supporting you

As long as you're taking WAINUA,
you're never alone.



Savings options

If your healthcare provider has prescribed WAINUA, you may have questions. AstraZeneca Access 360™ can discuss insurance, out-of-pocket costs, and financial assistance programs. Call 1-844-2-WAINUA (1-844-292-4682) to get started.



A dedicated point of contact

Connect with your Patient Education Manager throughout your treatment for:

- Information about WAINUA and hATTR-PN
- Updates during the insurance approval process
- Resources on how to take WAINUA



Educational resources

You'll receive information about how WAINUA works, education on hATTR-PN, and more.

*Among patients who have initiated WAINUA based on shipment data from Orsini.

†Current as of June 2024.

To Learn More or to Enroll in WAINUA WAY

Visit WAINUAWay.com or call
1-844-2-WAINUA (1-844-292-4682).



Actor portrayals.

You're in charge. We're here to help.

More than

68%

of patients are able to start
WAINUA within about

2 ^{*†}
WEEKS

Over

75%

of patients pay

\$0

out of pocket, with
savings programs that
help lower cost[†]



Tell your story Inspire people like you

The AZ US Patient Voices program is an opportunity to share your unique experiences with WAINUA. Email us at MyStory@AstraZeneca.com to learn more.

 **WAINUA**[®]
(eplontersen)^{45 mg}
injection for subcutaneous use



Actor portrayals.

SHINE A LIGHT ON AN hATTR-PN DIAGNOSIS

**Ask your healthcare provider
if WAINUA may be the right
treatment for you.**

Visit [WAINUA.com/Results](https://www.wainua.com/Results).

Please see Important Safety Information
on pages 4 and 5.

AstraZeneca  **IONIS**[®]

WAINUA is a registered trademark of the AstraZeneca group of companies.
©2025 AstraZeneca. All rights reserved. US-104731 Last Updated 12/25